



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Two new spermatophytes from California

ERNEST A. MCGREGOR

***Oxytheca Abramsii* sp. nov.**

Acaulescent or nearly so, 7–18 cm. high or less, sparingly dichotomously branched: stems glabrous mostly, but glandular just above the axils, herbaceous, becoming later more wiry: leaves spatulate, gradually narrowed to a wide petiole, 1.25–3 cm. long, scabrous-margined and the midvein slightly scabrous beneath; bracts ternate, slightly unequal, mucronate, ovate-lanceolate, connate and deflected mainly to one side, 3 mm. long: involucre broadly obconical, on peduncles 1–1.5 cm. long, those from the main axils rarely 2.5 cm. long; involucre-tube 1.5–2 mm. high; awns 10 (rarely 12), very fine, weak, but straight, 4 mm. long or less, the interval between the involucre ribs much wider than the ribs: flowers 6–8, at length nearly equaling the awns; pedicels equaling the perianth, well exserted from the involucre-tube: perianth about 4 mm. broad, cleft to base, lobes equal, elliptical, 2 mm. long, white or pinkish with wine-red midvein, slightly glandular and scabrous beneath: stamens inserted on perianth at base, shorter than the segments: styles 3, capitate: ovary glabrous.

O. Abramsii is nearest to *O. Parishii*, from which it differs in the following characters:

O. Parishii: Involucre ribs very prominent, firm, with the intervals almost lacking; awns 18–20, 3 times length of tube, strong, stiff, wiry; umbels 5–14-flowered; stems pale-glaucous, wiry.

O. Abramsii: Involucre ribs barely discernible, with intervals much wider than ribs; awns usually 10, not greatly exceeding the tube, inclined to be weak and herbaceous, though straight; umbels 6–8-flowered; stems green or reddish, not wiry.

Topatopa Mountains, Ventura Co., California. Growing on slopes in loose shale, elev. 1700 m. Collected by Abrams & McGregor (no. 72), June 5, 1908.

***Malacothrix arachnoidea* sp. nov.**

Perennial herb, with stout, freely branching, and very leafy herbaceous stem, 4.5–6 dm. high; herbage hoary throughout with

a dense woolly pubescence : leaves broadly lanceolate to linear-lanceolate, attenuate above, apiculate, tapering gradually below but sessile, and never at all auriculate or clasping, entire, or the



FIGURE 1. *Oxytheca Abramsii* McGregor. Natural size.

lower rarely remotely and obscurely dentate, 4–9 cm. long, reduced upward: heads terminating numerous branches, many-flowered; involucre 13–15 mm. high, hemispherical; bracts in about two series, becoming somewhat glabrate, the inner linear-lanceolate, acuminate, the outer similar but narrower and half as long; calyculate scales few: corolla white or pinkish; ligule 10 mm. long; tube 6 mm. long, puberulent on its upper third: achenes (not matured) broadly obovate-oblong, apparently 10-ribbed, the slightly contracted summit bearing a well-expanded crown whose rim is merely crenulate; pappus-bristles snow-white, all falling together.

Carmel Valley, Monterey Co., California; by the roadside. Collected by the writer (*no.* 1575), July 1, 1906.




FIGURE 2. *Oxytheca Abramsii* McGregor; a, leaf, natural size; b, bract, from free side, $\times 6$; c, bract, from deflected side, $\times 6$; d, involucre, showing ribs, awns, and flowers, $\times 5$; e, flower laid open to show parts, $\times 8$.

This plant belongs to the *M. saxatilis* group, which includes *M. saxatilis* (Nutt.) Torr. & Gray, *M. tenuifolia* (Nutt.) Torr. & Gray, *M. altissima* Greene, and *M. implicata* Eastw.

On the whole these species are very closely related and form a very natural group, differing from the other members of the genus in the following characters: tall, leafy perennials; involucre scales in two appressed equal series; receptacle with no bristles; no persistent pappus-bristles. It would seem that they might well constitute at least a valid subgenus.

The characters of the flower and those of the achene of the different species are much alike, the important differences being



FIGURE 3. *Malacothrix arachnoidea* McGregor. One half natural size.

in the aspect of the plants and the form of the leaves which, for a given type, seem quite distinctive.

Malacothrix arachnoidea is nearest to *M. saxatilis*, from which it is to be distinguished as follows :

M. saxatilis : Green, often woolly on young parts ; leaves obtuse, lower occasionally pinnatifid, auriculate, and partly clasping toward the base ; calyculate scales very numerous ; achenes crowned with a denticulate border.

M. arachnoidea : Hoary throughout with a dense woolly pubescence ; leaves apiculate-acute, apparently never pinnatifid, auriculate, or clasping ; the well-expanded crown of the achene with a merely crenulate border.

LELAND STANFORD JUNIOR UNIVERSITY.